

D-Brane Solutions in pp-wave Background

Alok Kumar, Rashmi R. Nayak and Sanjay

*Institute of Physics
Bhubaneswar 751 005
India*

email: kumar, rashmi, sanjay@iopb.res.in

Abstract

We present classical solutions for a $D5$ and $NS5$ -branes in a pp-wave background. The worldvolume coordinates for these branes lie along a six dimensional pp-wave configuration obtained from $AdS_3 \times S_3$ in a Penrose limit. One in addition has nontrivial R-R 3-form field strength as well as a dilaton background. Classical solutions for Dp -branes ($p > 5$), as well as their bound states, are then generated in both IIA and IIB string theories by using T -duality symmetries. We also briefly discuss the construction of Dp -branes in pp-wave backgrounds, obtained from string theories on $AdS_m \times S^m$. Finally, the construction of the above D5-brane solution is presented from the point of view of the Green-Schwarz worldsheet string theory and its low lying spectrum is discussed.

String theory in the presence of pp-wave backgrounds [1–7] has generated lots of excitement among physicists searching for a deep connection between quantum gravity and gauge theory dynamics. Such string backgrounds are technically tractable, and have direct applications to the four dimensional conformal theories from the point of view of a duality between string and gauge theories. The fact that pp-waves provide exact solutions of string theory, including in the presence of $R - R$ fields [4], is now known for long time. It has now been realized [8, 9] that in some special cases, they give rise to exactly solvable string theory with a quadratic action. As a result the spectrum of the theory can be explicitly obtained. Moreover, an interesting connection of pp-wave background with gauge field theories emerges by realizing that they can be obtained [10] in Penrose [11] scaling limit from a geometry of $AdS_p \times S^q$ type. This finally leads to a duality between string theories in pp-wave geometries and gauge theories, originally living on the boundary of the AdS spaces [12–16].

In view of the importance of BPS branes (preserving certain amount of supersymmetries) in understanding the non-perturbative dynamics of string theories, in this paper, we first present classical solutions of a $D5$ and an $NS5$ -brane in the background of a pp-wave. These are obtained from known classical solutions for such branes in $AdS_3 \times S^3$ background [17]. By setting the brane charges to zero, in our case, we obtain a ten dimensional solution with a six dimensional pp-wave background, giving rise to a massive Green-Schwarz string theory. Four spatial dimensions transverse to the branes, for these solutions, are flat.

To obtain our solution, we now begin by writing the $1_{NS} + 5_{NS} + 5'_{NS}$ solution given in [17]:

$$ds^2 = g_1^{-1}(x, y)(-dt^2 + dz^2) + H_5(x)dx^n dx^n + H'_5(y)dy^m dy^m, \quad (1)$$

$$dB = dg_1^{-1} \wedge dt \wedge dz + *dH_5 + *dH'_5, \quad (2)$$

$$e^{2\phi} = \frac{H_5(x)H'_5(y)}{g_1(x, y)}, \quad (3)$$

where

$$[H'_5(y)\partial_x^2 + H_5(x)\partial_y^2]g_1(x, y) = 0. \quad (4)$$

A particular solution for g_1 is given as:

$$g_1(x, y) = H_1(x)H'_1(y), \quad (5)$$

with

$$H_1 = 1 + \frac{R_1^2}{x^2}, \quad H_5 = 1 + \frac{R_5^2}{x^2}, \quad H'_1 = 1 + \frac{R_1'^2}{y^2}, \quad H'_5 = 1 + \frac{R_5'^2}{y^2}. \quad (6)$$

By setting $H'_1 = 1$, the above solution becomes a direct product of NS1 + NS5 configuration, together with an NS5-brane. Then, in a decoupling limit $x \rightarrow 0$, one obtains from the metric given in (1) [17]:

$$ds^2 = \frac{x^2}{R_1^2}(-dt^2 + dz^2) + R_1^2 \frac{dx^2}{x^2} + R_1^2 d\Omega_3^2 + H'_5(y)(dy^2 + y^2 d\Omega_3'^2), \quad (7)$$

where one has also used $H_1 = H_5$ to simplify the expression. Finally, by using an S-duality transformation on the background mentioned above, one obtains a metric:

$$ds^2 = H_5'^{-1/2} \left[\frac{x^2}{R_1^2}(-dt^2 + dz^2) + R_1^2 \frac{dx^2}{x^2} + R_1^2 d\Omega_3^2 \right] + H_5'^{1/2}(y)(dy^2 + y^2 d\Omega_3'^2). \quad (8)$$

We now obtain the $D5$ brane solution in the pp-wave background, by interpreting the metric in eqn.(8) as that of a $D5$ -brane in $AdS_3 \times S_3 \times R^4$. The coordinates t, x and z in the above solution represent the AdS_3 space, whereas $d\Omega_3^2$ is the metric of an S^3 . It is also evident that the worldvolume of the above $D5$ -brane solution coincides with this $AdS_3 \times S^3$ space and transverse directions to the branes are represented by the coordinates y^m . We now apply the Penrose limit [10] on t, x, z , as well as on the coordinates of S^3 parametrized as:

$$d\Omega_3^2 = d\psi^2 + \sin^2\psi d\Omega_2^2, \quad (9)$$

on the $D5$ -brane metric in (8). Then, by following the procedure in [10], one obtains from the metric in (8):

$$dS^2 = H_5'^{-1/2} [dudv - \mu^2 \sum_{i=1}^4 z^{i2} du^2 + \sum_{i=1}^4 dz^i dz^i] + H_5'^{1/2} (dy^2 + y^2 d\Omega_3'^2). \quad (10)$$

As is noticed, the y dependence in the limiting procedure described above remains unchanged due to a simultaneous scaling of y together with R_5' .

The expression for the dilaton field in the Penrose limit above can be found rather easily and is given by

$$e^{2\phi} = H_5'(y)^{-1}. \quad (11)$$

The expression for the $(R-R)$ (3rd rank antisymmetric) field strength (H^R) in the pp-wave limit is obtained from the NS-NS field strength given in expression (2) by applying S-duality transformation on these fields. As is noticed, in this case there are several nonzero components. In the pp-wave limit, the constant field strengths: H_{+12}^R and H_{+34}^R emerge from the first two terms in the RHS of eqn. (2), in two steps. First one obtains, in $x \rightarrow 0$ decoupling limit, expression for the field strength that is given by the appropriate levi-civita tensors on AdS_3 [18] and S^3 respectively. In the second step, one applies the Penrose limit following the procedure mentioned above to get constant field strengths [12]:

$$H_{+12}^R = H_{+34}^R = 2\mu. \quad (12)$$

We however notice that the last term in the RHS of eqn. (2) also survives the above limiting processes and its functional form remains unchanged. As a result we have additional nonzero components of the $R - R$ 3-form field strength [19]:

$$H_{mnp}^R = \epsilon_{mnp\ell} \partial_\ell H_5'(y). \quad (13)$$

Here we once again like to emphasize that all the three terms: representing the field strengths on AdS_3 , S^3 and R^4 , of a $D5$ brane solution, have identical scaling properties, when standard Penrose scalings of [10] are applied in the $AdS_3 \times S^3$ part and transverse

coordinates are scaled appropriately. We have therefore presented a classical solution corresponding to a $D5$ -brane in an exactly solvable pp-wave background.

We can also write down the NS5 brane solution in a pp-wave background using the metric in equation (7). One now obtains the metric:

$$dS^2 = [dudv - \mu^2 \sum_{i=1}^4 z^{i2} du^2 + \sum_{i=1}^4 dz^i dz^i] + H'_5(dy^2 + y^2 d\Omega_3^2). \quad (14)$$

The dilaton now has a form:

$$e^{2\phi} = H'_5(y). \quad (15)$$

The antisymmetric tensors are exactly of the form in eqns. (12) and (13), with ‘R’ replaced by ‘NS’ in these equations.

Now we generate Dp -branes ($p > 5$), starting with our solution as given in eqns.(10)-(13) by applying T-duality [20, 21] along the transverse directions of the $D5$ -brane. Applying this procedure to a transverse coordinate $\tilde{x}$ one obtains a $D6$ brane in type IIA theory. The explicit solution for the non-zero components of the metric, field strengths and dilaton are then given as follows:

$$\begin{aligned} e^{2\phi_a} &= H_6^{-3/2}, \\ A_m^{(1)} &= A_{\tilde{x}m}^{(R)} : F_{mn}^{(1)} = \epsilon_{mnp} \partial_p H_6, \\ A_{\tilde{x}\mu\nu}^{(3)} &= A_{\mu\nu}^{(R)}, \quad \tilde{F}_{\tilde{x}\alpha\beta\gamma} = \partial_\gamma A_{\tilde{x}\alpha\beta} + cycl. = H_{\alpha\beta\gamma}^{(R)} : \quad \tilde{F}_{\tilde{x}u12} = \tilde{F}_{\tilde{x}u34} = 2\mu, \end{aligned} \quad (16)$$

$$dS^2 = H_6^{-1/2} [dudv - \mu^2 \sum_{i=1}^4 z^{i2} du^2 + \sum_{i=1}^4 dz^i dz^i + d\tilde{x}^2] + H_6^{1/2} (dr^2 + r^2 d\Omega_2^2), \quad (17)$$

with $H_6 = (1 + R_6/r)$, r being the radial coordinate in transverse direction and $H^{(R)}$ denotes the field strength that we found earlier for the $D5$ brane solution in the IIB theory. Integer subscripts in field strenghts above denote coordinates z^i 's and indices μ, ν run over the longitudinal directions. For $\mu = 0$ this solution reduces to the standard

$D6$ brane in type IIA theories [22]. One can continue this exercise further to the higher branes as well. Then, for example, a logarithmic Green function appears for a $D7$ brane and a linear function appears in the solution for a $D8$ brane [23]. The explicit solution for the $D7$ brane ($H_7 = 1 - \mu \ln(r/l)$) [21] is given as:

$$\begin{aligned}
e^{2\phi_b} &= H_7^{-2}, \\
\chi &= -A_{\tilde{y}}^{(1)} : \quad \partial_m \chi = -\epsilon_{mn} \partial_n H_7, \\
F_{\tilde{x}\tilde{y}u12}^{(5)} &= F_{\tilde{x}\tilde{y}u34}^{(5)} = 2\mu, \\
dS^2 &= H_7^{-1/2} [dudv - \mu^2 \sum_{i=1}^4 z^{i2} du^2 + \sum_{i=1}^4 dz^i dz^i + d\tilde{x}^2 + d\tilde{y}^2] + H_7^{1/2} (dr^2 + r^2 d\theta^2). \quad (18)
\end{aligned}$$

Solutions corresponding to the bound states of Dp branes can also be constructed using the procedure given in [21]. For example, a $D5 - D7$ bound state solution can be obtained from the $D6$ brane solution in eqns.(16) and (17) by applying a rotation between the longitudinal coordinate $\tilde{x}$ and one of the transverse coordinate $\tilde{y}$ for this solution, and then applying T-duality along (rotated) coordinate $\tilde{y}'$. We skip the details of this solution and now briefly discuss the construction of Dp branes in other pp-wave backgrounds. These backgrounds can be obtained from string theory on $AdS_m \times S^m$ for some other values of m . In $m = 2$ case, for example, relevant field strength living on AdS_2 and S^2 now correspond to a 1-form field. It should therefore be possible to construct $D4$ branes in a pp-wave background, now originating from $AdS_2 \times S^2 \times R$. One now has R^5 as a five dimensional transverse space on which a 4-form field strength of the type: $F_{mnpq} = \epsilon_{mnpqr} \partial_r H_4$ lives. One in addition will have constant field strengths of the type: $F_{+1}^{(1)} = F_{+2}^{(1)} = \mu$. Once again, by applying T -dualities, all the higher branes can be obtained. This process can also be repeated for $D8$ branes in a pp-wave background obtained from $AdS_4 \times S^4 \times R$. We hope to come back with explicit solutions of these types in near future.

We now proceed to construct the $D5$ -brane, obtained earlier as a classical solution in supergravity, from the point of view of a first quantized string theory in Green-Schwarz

formalism in light-cone gauge. The relevant classical action to study in our case, obtained from $AdS_3 \times S^3 \times R^4$ by applying the Penrose limit, has a form [9]:

$$L = L_B + L_F,$$

$$L_B = \partial_+ u \partial_- v - m^2 x_i^2 + \partial_+ x_i \partial_- x_i + \partial_+ x_\alpha \partial_- x_\alpha. \quad (20)$$

$$L_F = i\theta_R \gamma^v \partial_+ \theta_R + i\theta_L \gamma^v \partial_- \theta_L - 2im\theta_L \gamma^v M \theta_R, \quad (21)$$

$$m \equiv \alpha' p^u \mu = 2\alpha' p_v \mu, \quad (22)$$

$$M = -\frac{1}{2}(\gamma^{12} + \gamma^{34}), \quad (23)$$

where (θ_R, θ_L) are 16-component Majorana-Weyl spinors in the right and left moving sectors. x_i and x_α in eqn.(20) refer to the longitudinal $D5$ brane coordinates z^i and transverse ones y^m in the metric given in eqn.(1). The form of the matrix M in the present case leads to only half of the fermions picking up masses [9]. This is different from the case of pp-wave background that is obtained from $AdS_5 \times S^5$ [24, 25], and leads to some differences in our case compared to the one in [24].

The equations of motion following from (20) has a form:

$$\partial_+ \partial_- x_i + m^2 x_i = 0, \quad \partial_+ \partial_- x_\alpha = 0, \quad (24)$$

$$\partial_+ \theta_R - mM\theta_L = 0, \quad \partial_- \theta_L - mM\theta_R = 0. \quad (25)$$

After solving the light-cone gauge condition one rewrites the above action, as well as equations of motion using 8-component spinors S_L and S_R . The equations of motion have an identical form as in (24), (25) by replacing $\theta_{L,R}$ by $S_{L,R}$. The matrix M now is an 8×8 one of the same structure as in (23). Keeping in mind the explicit form of the matrix M , which evidently breaks the $SO(8)$ further, one splits the fermions in the $8 \rightarrow 4 + 4$ way: $S_L \rightarrow (\tilde{S}_L, \hat{S}_L)$, $S_R \rightarrow (\tilde{S}_R, \hat{S}_R)$:

$$\gamma^{1234} \begin{pmatrix} \tilde{S}_{L,R} \\ \hat{S}_{L,R} \end{pmatrix} = \begin{pmatrix} -\tilde{S}_{L,R} \\ \hat{S}_{L,R} \end{pmatrix}. \quad (26)$$

Finally one introduces 4×4 matrices Λ and Σ :

$$\gamma^{12} \begin{pmatrix} \tilde{S}_{L,R} \\ \hat{S}_{L,R} \end{pmatrix} = - \begin{pmatrix} \Lambda \tilde{S}_{L,R} \\ \Sigma \hat{S}_{L,R} \end{pmatrix}, \quad (27)$$

with $\Lambda^2 = \Sigma^2 = -1$. Λ and Σ in the above equation are 4×4 antisymmetric matrices with eigenvalue $\pm i$. Using these notations one has:

$$M \begin{pmatrix} \tilde{S}_{L,R} \\ \hat{S}_{L,R} \end{pmatrix} = \begin{pmatrix} \Lambda \tilde{S}_{L,R} \\ 0 \end{pmatrix}. \quad (28)$$

The equations of motion written in terms of $(\tilde{S}_L, \hat{S}_L)$ and $(\tilde{S}_R, \hat{S}_R)$ are then of the form:

$$\partial_+ \tilde{S}_R - m \Lambda \tilde{S}_L = 0, \quad \partial_- \tilde{S}_L - m \Lambda \tilde{S}_R = 0, \quad (29)$$

$$\partial_+ \hat{S}_R = 0, \quad \partial_- \hat{S}_L = 0. \quad (30)$$

We now write down the open string boundary condition [24] corresponding to the above $D5$ -brane configuration. These are given on the longitudinal bosonic coordinates x^i , ($i = 1, \dots, 4$) and transverse ones x^α ($\alpha = 5, \dots, 8$) by standard Neumann and Dirichlet conditions. For spinors we have:

$$S_L|_{\sigma=0,\pi} = \Omega S_R|_{\sigma=0,\pi}, \quad (31)$$

with Ω being the product of γ matrices along the Dirichlet directions. Ω has to satisfy the following consistency conditions in the present case:

$$[\Omega, \gamma] = 0, \quad (32)$$

$$\Omega M \Omega = M. \quad (33)$$

The first of these equations follows from the requirement that S_L and S_R have the same $SO(8)$ chirality. The second condition is a consequence of a consistency condition on the above equations of motion in the zero mode sector, leading to one half supersymmetry.

The explicit form of Ω in our case is given by $\Omega = \gamma^{5678}$ and leads to the following boundary conditions on $(\tilde{S}_L, \hat{S}_L)$ and $(\tilde{S}_R, \hat{S}_R)$ using eqn.(26):

$$\tilde{S}_L|_{\sigma=0,\pi} = -\tilde{S}_R|_{\sigma=0,\pi}, \quad (34)$$

$$\hat{S}_L|_{\sigma=0,\pi} = \hat{S}_R|_{\sigma=0,\pi}. \quad (35)$$

Now, as the equations of motion as well as the boundary condition for the components $\hat{S}_{L,R}$ are identical to the ones in flat space, we only concentrate on finding explicit solution for $\tilde{S}_{L,R}$ below.

Explicit solution for x^i and x^α are given as:

$$x^i(\sigma, \tau) = [x_0^i \cos m\tau + \frac{1}{m} p_0^i \sin m\tau] + i \sum_{n \neq 0} \frac{1}{\omega_n} \alpha_n^i e^{-i\omega_n \tau} \cos n\sigma, \quad (36)$$

$$x^\alpha(\sigma, \tau) = \sum_{n \neq 0} \frac{1}{n} \alpha_n^\alpha e^{-in\tau} \sin n\sigma, \quad (37)$$

where $\omega_n = \text{sgn}(n)\sqrt{n^2 + m^2}$ and oscillators satisfy the following commutation relations:

$$[a_n^I, a_l^J] = \delta_{n+l} \delta^{IJ}, \quad [\bar{a}_0^i, a_0^j] = \delta^{ij}, \quad (38)$$

with indices I, J running over all the eight coordinates. We have also used the definitions:

$$a_0^i = \frac{1}{\sqrt{2m}}(p_0^i + imx_0^i), \quad \bar{a}_0^i = \frac{1}{\sqrt{2m}}(p_0^i - imx_0^i), \quad a_n^I = \sqrt{\frac{1}{|\omega_n|}} \alpha_n^I. \quad (39)$$

For $\tilde{S}_{L,R}$ we have the solution of the equations of motion [9]:

$$\tilde{S}_L = [\tilde{S}_0^a \cos \omega_0 \tau - \tilde{S}_0^b \sin \omega_0 \tau] + \sum_{n \neq 0} \frac{c_n}{\sqrt{2}} [\tilde{\phi}_n \tilde{S}_n^a + \frac{i}{m} (\omega_n - n) \Lambda_b^a S_n^b \phi_n], \quad (40)$$

$$\tilde{S}_R = \Lambda_b^a [\tilde{S}_0^b \cos \omega_0 \tau + \tilde{S}_0^a \sin \omega_0 \tau] + \sum_{n \neq 0} \frac{c_n}{\sqrt{2}} [\phi_n S_n^a + \frac{i}{m} (\omega_n - n) \Lambda_b^a \tilde{S}_n^b \tilde{\phi}_n], \quad (41)$$

where $c_n = m/\sqrt{m^2 + (\omega_n - n)^2}$ and

$$\tilde{\phi}_n = \exp -i(\omega_n \tau + n\sigma), \quad \phi_n = \exp -i(\omega_n \tau - n\sigma), \quad (S_{-n} = S_n^\dagger \text{ etc.}) \quad (42)$$

Moreover, the boundary condition eqn. (31) implies:

$$\tilde{S}_0^a = -(\Lambda \tilde{\tilde{S}}_0)^a, \quad \tilde{S}_n^a = -S_n^a. \quad (43)$$

One can now put the conditions (43) back into the solutions in eqns. (40) and (41) to find out that indeed the boundary conditions in eqns. (34) and (35) are being satisfied. The commutation relations for the above spinor components turn out to be of the form:

$$\{\tilde{S}_0^a, \tilde{S}_0^b\} = \delta^{ab}, \quad \{S_n^a, S_{-n}^b\} = \delta^{ab}. \quad (44)$$

The total Hamitlonian can be written as:

$$H = H_0 + \tilde{H} + \hat{H}, \quad (45)$$

where

$$\begin{aligned} H_0 &= \frac{m}{p^u} (\bar{a}_0^i a_0^i - i \tilde{S}_0^a \Lambda \tilde{\tilde{S}}_0^a) + e_0, \\ \tilde{H} &= \frac{1}{p^u} \sum_{n=1}^{\infty} \omega_n (a_n^{i\dagger} a_n^i + S_n^{a\dagger} S_n^a), \\ \hat{H} &= \frac{1}{p^u} \sum_{n=1}^{\infty} n (a_n^{\alpha\dagger} a_n^\alpha + \hat{S}_n^{A\dagger} \hat{S}_n^A). \end{aligned} \quad (46)$$

with e_0 being the zero point energy. The form of the zero mode part of the hamiltonian implies that the ground state degeneracy of a $D5$ -brane spectrum is broken. To analyze the spectrum of the states, we use the fermionic creation and annihilation operators constructed out of spinors: $\tilde{S}$ and $\hat{S}$. Since both of these are 4 component objects, each of them give rise to two such creation operators and two annihilation opertaors. Then sixteen states (at a level defined by the vacuum energy $e_0 = 2\mu$ [24]), in this representation, are obtained using these operators on appropriate Fock vacuum for $\mu = 0$. On the other hand for $\mu \neq 0$, there is a splitting in this spectrum which can now be analyzed by simply taking into account the H_0 part in eqns.(45) and (46) .

We have therefore obtained explicit construction of the open string describing the $D5$ -branes in the pp-wave background presented earlier in this paper. It will be interesting to generalize these results to other D-branes and their bound states, discussed in the paper, as well. We however leave this as a future exercise.

References

- [1] R.Gueven, Phys.Lett. **B 191** (1987) 275.
- [2] D. Amati, C. Klimcik, Phys.Lett. **B 210** (1988) 92.
- [3] G. T. Horowitz, A. A. Tseytlin, Phys. Rev **D 51** (1995) 2896, [hep-th/9409021].
- [4] S. Kar, A. Kumar, G. Sengupta, Phys. Lett. **B 375** (1996) 121, [hep-th/9601171].
- [5] A. Kumar and G. Sengupta, Phys.Rev. **D54** (1996) 3976, [hep-th/9602145].
- [6] G. Sengupta, Phys.Rev. **D55** (1997) 3793, [hep-th/ 9609152]
- [7] R. Gueven, Phys. Lett. **B482** (2000) 255, [hep-th/0005061].
- [8] R.R Metsaev, Nucl.Phys.B **B625** (2002) 70, [hep-th/0112044].
- [9] J.G.Russo, A.A.Tseylin, [hep-th/0202179].
- [10] Matthias Blau, Jos Figueroa-O’Farrill, George Papadopoulos, [hep-th/0202111].
- [11] R. Penrose, "Any space-time has a plane wave as a limit", *in Differential geometry and Relativity*, pp. 271-275, Reidel, Dordrecht, (1976).
- [12] D. Berenstein, J. Maldacena, H. Nastase , [hep-th/0202021].
- [13] N. Itzhaki, Igor R. Klebanov, Sunil Mukhi, [hep-th/0202153].
- [14] L. Zayas, J. Sonnenschein, [hep-th/0202186].
- [15] M. Alishahiha, M.M. Sheikh-Jabbari, [hep-th/0203018]
- [16] U. Gursoy, C. Nunez and M. Schwelling, [hep-th/0203124].
- [17] G. Papadopoulos, J. Russo, A.A. Tseytlin, Class. Quant. Grav. **17** (2000)1713, [hep-th/9911253].

- [18] A.Ali, A.Kumar,Mod.Phys.Lett.**A8** (1993) 2045 [hep-th/9303032].
- [19] A.A. Tseytlin, Mod.Phys.Lett. **A11** (1996) 689, [hep-th/9601177].
- [20] E. Bergshoeff, C. Hull, T. Ortin, Nucl. Phys. **B451** (1995) 547,[hep-th/9504081].
- [21] J. C. Breckenridge, G. Michaud, R. C. Myers, Phys. Rev. **D55** (1997) 6438, [hep-th/9611174].
- [22] M.J. Duff, Ramzi R. Khuri and J.X. Lu, Phys.Rept.**259** (1995) 213, [hep-th/9412184].
- [23] E. Bergshoeff, M. de Roo, M.B. Green, G. Papadopoulos and P.K. Townsend, Nucl.Phys. **B470** (1996) 113, [hep-th/9601150].
- [24] A. Dabholkar, S. Parvizi, [hep-th/0203231].
- [25] see also P. Lee and J. Park, [hep-th/0203257].